Turkish Journal of Zoology

Volume 38 | Number 5 Article 8

1-1-2014

# Development of empirical standard weight equation for Pursak chub Squalius pursakensis, an endemic cyprinid species of Northwest Anatolia

ŞULE SÜLÜN

SERCAN BAŞKURT

ÖZGÜR EMİROĞLU

DANIELA GIANNETTO

ALİ SERHAN TARKAN

*See next page for additional authors*

Follow this and additional works at: https://journals.tubitak.gov.tr/zoology

 Part of the Zoology Commons

## Recommended Citation

SÜLÜN, ŞULE; BAŞKURT, SERCAN; EMİROĞLU, ÖZGÜR; GIANNETTO, DANIELA; TARKAN, ALİ SERHAN; AĞDAMAR, SEVAN; GAYGUSUZ, ÖZCAN; DORAK, ZEYNEP; AYDIN, HAMDİ; and ÇİÇEK, ARZU (2014) "Development of empirical standard weight equation for Pursak chub Squalius pursakensis, an endemic cyprinid species of Northwest Anatolia," *Turkish Journal of Zoology*: Vol. 38: No. 5, Article 8.
https://doi.org/10.3906/zoo-1311-24
Available at: https://journals.tubitak.gov.tr/zoology/vol38/iss5/8

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Zoology by an authorized editor of TÜBİTAK Academic Journals. For more information, please contact academic.publications@tubitak.gov.tr.

# Development of empirical standard weight equation for Pursak chub Squalius pursakensis, an endemic cyprinid species of Northwest Anatolia

Authors
ŞULE SÜLÜN, SERCAN BAŞKURT, ÖZGÜR EMİROĞLU, DANIELA GIANNETTO, ALİ SERHAN TARKAN, SEVAN AĞDAMAR, ÖZCAN GAYGUSUZ, ZEYNEP DORAK, HAMDİ AYDIN, and ARZU ÇİÇEK

This article is available in Turkish Journal of Zoology: https://journals.tubitak.gov.tr/zoology/vol38/iss5/8

TÜBİTAK

**Turkish Journal of Zoology**

**http://journals.tubitak.gov.tr/zoology/**

**Research Article**

Turk J Zool
(2014) 38: 582-589
© TÜBİTAK
doi:10.3906/zoo--1311-24

# Development of empirical standard weight equation for Pursak chub *Squalius pursakensis*, an endemic cyprinid species of Northwest Anatolia

**Şule SÜLÜN[1], Sercan BAŞKURT[1], Özgür EMİROĞLU[1], Daniela GIANNETTO[2,*], Ali Serhan TARKAN[2], Sevan AĞDAMAR[2], Özcan GAYGUSUZ[3], Zeynep DORAK[3], Hamdi AYDIN[4], Arzu ÇİÇEK[5]**

[1]Department of Biology, Faculty of Science, Eskişehir Osmangazi University, Eskişehir, Turkey
[2]Faculty of Fisheries, Muğla Sıtkı Koçman University, Kötekli, Muğla, Turkey
[3]Faculty of Fisheries, İstanbul University, Laleli, Fatih, İstanbul, Turkey
[4]Gazanfer Bilge Vocational High School, Kocaeli University, Karamürsel, Kocaeli, Turkey
[5]Environmental Problems Research and Application Center, Anadolu University, Eskişehir, Turkey

**Received:** 11.11.2013 • **Accepted:** 10.03.2014 • **Published Online:** 14.07.2014 • **Printed:** 13.08.2014

**Abstract:** Indices of condition enable the evaluation of the well-being of fish, with the assumption that heavier fish of a given length are in a better condition. Relative weight ($W_r$) is one of these indices; it is calculated by comparing the actual weight of a specimen with the ideal weight of a specimen in good physiological condition of the same length from the same species, i.e. standard weight ($W_s$). In this research, length and weight data over the distribution range for Pursak chub *Squalius pursakensis*, an endemic species distributed in the Sakarya and Porsuk drainages in Northwest Anatolia (Turkey), were used to compute a $W_s$ equation by means of the empirical percentile (EmP) method. The $W_s$ equation obtained was $\log_{10} W_s = -4.657 + 2.614 \log_{10} TL + 0.127 (\log_{10} TL)^2$, and the total length range of application was 80–340 mm. Since the EmP $W_s$ equation was not influenced by length variation, the use of this equation to compute the relative weight ($W_r$) for *S. pursakensis* throughout its area of distribution is suggested.

**Key words:** Condition indices, endemic species, *Squalius*, relative weight

## 1. Introduction

Indices of condition are commonly used by fisheries to evaluate the well-being of fish, starting with the assumption that "fatter is fitter" (sensu Glazier, 2000). Thus, comparing specimens of the same length, fish with a higher weight are in better condition than those with a lower weight (Anderson and Neumann, 1996; Blackwell et al., 2000). These indices are important tools for fishery management (Anderson and Neumann, 1996; Blackwell et al., 2000) and have been used in fishery research since the beginning of the 20th century (Froese, 2006). According to Fechhelm et al. (1995), the use of indices of condition to monitor the well-being of fish is cost-effective because, being based only on length and weight measurements, indices do not require the sacrifice of the specimens and allow evaluation of the condition of fish with minimal mortality.

Relative weight ($W_r$) (Wege and Anderson, 1978) is one of these indices and, compared to the others available in literature [i.e. Fulton's (1911) condition factor and Le Cren's (1951) condition factor], it has the great advantages of not being influenced by changes in body shape and of allowing for comparing the condition of fish of different lengths belonging to different populations (Murphy et al., 1991). Relative weight is based on the comparison between the actual weight of a specimen and the standard weight ($W_s$), which is the weight in the same length of an ideal fish of the same species that is in good physiological condition (Murphy et al., 1991). $W_s$ is predicted by a standard weight equation, that is, a length–weight regression typical of the species (Wege and Anderson, 1978). Since its development, the $W_r$ index has been widely used to perform condition analyses of many species (Blackwell et al., 2000). However, its applicability is limited by the lack of species-specific standard weight equations that have to be developed using a wide sample of specimens collected throughout the area of distribution of each species (Bister et al., 2000).

Pursak chub *Squalius pursakensis* (Hankó, 1925) is a cyprinid species endemic to the Sakarya and Porsuk drainages (Turkey), which flow into the Black Sea (Northwest Anatolia) (Özulug and Freyhof, 2011) (Figure 1). The species, originally described as *Leuciscus orientalis pursakensis* Hankó, 1925, was recently rehabilitated as a valid species by Özulug and Freyhof (2011). On the basis of molecular differences found in chubs from West

* Correspondence: danielagiannetto@gmail.com

Anatolia, Durand et al. (2000) suggested that the group of "short snouted chubs", previously identified as a subspecies of *Squalius cephalus* (Linnaeus, 1758), could be divided into different species. Recently, Özulug and Freyhof (2011) confirmed this hypothesis and identified the presence of 10 different species of the genus *Squalius* in West and Central Anatolia (4 of them only recently described), including *S. pursakensis*.

This species is consumed by local people of the basin and is used for recreational purposes (Erk'akan, 1981; Ekmekçi, 1996; Emiroğlu et al., 2001). As both basins (Sakarya and Porsuk) where *S. pursakensis* occurs have recently been severely polluted and are affected by serious habitat destruction, mainly damming and water abstraction, the species has suffered from these changes (Bostancı and Polat, 2009; Innal, 2010).

Two different methods have been proposed in the literature for the computation of $W_s$ equations: the regression line percentile (RLP) method, proposed by Murphy et al. (1990), and the empirical percentile (EmP) method, developed by Gerow et al. (2005). Until recently, the most widely used method was the RLP (Blackwell et al., 2000). However, according to Gerow et al. (2004), some of the $W_s$ equations developed using the RLP method showed significant length-related biases; moreover, the results of recent studies have encouraged the use of the EmP method in developing $W_s$ equations (Rennie and Verdon, 2008; Ogle and Winfield, 2009; Gerow, 2010; Giannetto et al., 2011, 2012a).

The aim of this research was to develop a standard weight equation by means of the EmP method for *S. pursakensis*, which could allow estimating the relative weight throughout the entire area of distribution of this species. Given that the species is under threat from habitat disturbance and biotic degradations by nonnative species, which are commonly increasing in the area, the proposed standard weight equation would serve as an important tool for conservation of this endemic species in its restricted distribution area.

## 2. Materials and methods

### 2.1. Dataset selection

Specimens of *S. pursakensis* were collected throughout the area of distribution of the species (Figure 2) by means of electrofishing (SAMUS 725G) between 2009 and 2012. After collection, each fish was measured for length [total length (TL), standard length (SL), and fork length (FL)] to the nearest 1 mm and wet weight (W) to the nearest 0.1 g in situ, with fish being returned to the water.

According to Giannetto et al. (2011, 2012c), the following methods were used to validate the total dataset. First, by the TL–W regression of the total sample, all specimens that were large outliers were excluded, as they were probably derived from incorrect measurements. The dataset was then separated into statistical populations on the basis of the data and location of collection (Ogle and Winfield, 2009). In this regard, a stock (population) was regarded as different according to sampling years and periods from the same location because of emigration/immigration, recruitment, and mortality (Begg and Waldman, 1999). Populations with fewer than 10 specimens were removed from the dataset (Ogle and Winfield, 2009; Lorenzoni et al., 2012). Hence, in order to identify all anomalous values, data were further validated by plotting the $\log_{10}$TL – $\log_{10}$W regression for each population separately; all values that diverged by more than double the expected value from the regression curve were removed (Bister et al., 2000). These equations were then analyzed, and all populations showing an $R^2$ value of less than 0.90 or for which the value of the slope (*b*) fell outside the range of 2.5–3.5 were excluded from further analyses (Froese, 2006).



**Figure 1**. *Squalius pursakensis* from Seydi Suyu in Sakarya Basin (445 mm TL).

The last step for the validation was to plot the slopes (*b*) of all populations against all intercepts ($\log_{10} a$) (Pope et al., 1995) to identify the outliers of this regression, represented by populations composed of a few fish or samples with a narrow length range (Froese, 2006).

### 2.2. Determination of the applicable total length range for the $W_s$ equation

Once the dataset was validated, the next step was to determine a suitable length range for the application of the $W_s$ equation. A minimum length is preferred because small fishes show a high variance due to the differences in growth forms that arise in the juvenile stages, and due to the potential for error associated with their measurement in the field (Murphy et al., 1990; Rypel and Richter, 2008). In accordance with Willis et al. (1991), the minimum total length was determined by plotting the variance/mean ratio for $\log_{10}$W on 10-mm total length intervals, and the minimum length was determined as the size at which the value of the ratio was less than 0.01 (Murphy et al., 1990).

Since the EmP method was utilized to develop the $W_s$ equation, a maximum length for the application of that equation was also required (Gerow et al., 2005). This length was determined as the length class for which at least




**Figure 2.** Area of distribution of *Squalius pursakensis* (dark gray area).

3 fish populations were present in the dataset, because 3 represents the smallest sample size that allows estimation of quartiles (Gerow et al., 2005). Any fish outside the suitable length range were not utilized for further analyses.

Next, the dataset, thus validated, was divided into 2 sets: a larger development dataset (used to compute the $W_s$ equation) and a smaller validation dataset (used to assess potential length-related biases in the $W_s$ equation) (Rypel and Richter, 2008; Ogle and Winfield, 2009; Giannetto et al., 2012c).

### 2.3. Development of the $W_s$ equation

The EmP method proposed by Gerow et al. (2005) was used to develop the $W_s$ equation for *S. pursakensis*. The mean empirical W for each 10-mm length-group was calculated by the $\log_{10}$-transformed TL and W of each population of the development dataset; the third quartiles of these mean empirical W values for each length-group were then regressed against TL to develop the EmP $W_s$ equation by using a weighted quadratic model (Gerow et al., 2005).

Using the $W_s$ equation obtained, the $W_r$ of each specimen from each population was then calculated by means of the equation provided by Wege and Anderson (1978): $W_r = 100\ (W / W_s)$, where W is the weight of an individual in grams and $W_s$ is the standard weight predicted by the $W_s$ equation.

### 2.4. Validation of the EmP $W_s$ equation

The principal attribute of a good condition index is that, in order to enable accurate comparisons of condition assessments of fish of different sizes, the measure of the condition should be free from length-related biases (Murphy et al., 1991; Anderson and Neumann, 1996; Blackwell et al., 2000). To this aim, 3 different methods were used to detect potential length-related biases and to validate the EmP $W_s$ equation calculated for *S. pursakensis*: the analysis of the residuals of the $W_s$ equation, to investigate whether the distribution of residuals exhibits evident patterns (Ogle and Winfield, 2009; Giannetto et al., 2012a); the Willis method (Willis et al., 1991), whereby a chi-square test is used to determine whether the proportion of populations with a significant positive slope in the TL–$W_r$ equation is equal to the proportion of those with a significant negative slope; and the empirical quartiles (EmpQ) method (Gerow et al., 2004) as modified by Ogle and Winfield (2009), applied to the validation dataset by using the FSA package (http://www.rforge.net/FSA) of R Software to determine if the slope of the quadratic regression of the third quartile of the mean W standardized by $W_s$ against length intervals of 10 mm had a value of zero (Ogle and Winfield, 2009; Giannetto et al., 2011, 2012a, 2012c).

## 3. Results

### 3.1. Dataset selection

A total of 2590 *S. pursakensis* specimens were collected during the research. The total length of specimens collected ranged from a minimum of 26.000 mm to a maximum of 445.000 mm (mean length: 152.562 ± 51.348 mm) and the weight from a minimum of 0.200 g to a maximum of 1263 g (mean weight: 62.174 ± 82.197 g). A new maximum total length (445 mm) for this species was recognized.

The log-transformed TL–W equation calculated for the total sample was:

$\log_{10}(W) = -5.304 + 3.176 \log_{10} TL$ (mm) ($R^2 = 0.985$, $P < 0.001$; n = 2590).

The SL–TL and FL–TL equations resulted as follows:

TL = 5.724 + 1.157 SL ($R^2$ = 0.997; n = 1684)

and

TL = 3.225 + 1.053 FL ($R^2$ = 0.998; n = 1684).

The total dataset was then divided into 78 statistical populations. By the linear plot between $\log_{10}(a) - b$, no populations were identified as outliers, and for all 78 the value of $R^2$ was >0.95 and the *b* value was within the range of 2.5–3.5.

### 3.2. Determination of the applicable total length range for the $W_s$ equation

The minimum TL for the application of the $W_s$ equation was determined as 80 mm (Figure 3). The maximum TL was assigned to be 340 mm because, despite the presence in the dataset of fish longer than this size, 340 mm was the length at which at least 3 values were present (Table). According to this, the length range judged to be suitable for the EmP $W_s$ equation was 80–340 mm.

The total sample was then divided into 2 datasets: a development dataset made up of 2210 specimens, and a validation dataset of 380 fish.

### 3.3. Development of $W_s$ equation

By applying the EmP method to the development dataset, the $W_s$ equation for *S. pursakensis* resulted as follows:

$\log_{10} W_s = -4.657 + 2.614 \log_{10} TL + 0.127 (\log_{10} TL)^2$ ($R^2$ = 0.999, P < 0.001).

### 3.4. Validation of the EmP $W_s$ equation

The residual values of the EmP $W_s$ equation showed a random distribution and did not exhibit evident patterns (Figure 4). Applying the Willis method, the number of TL–W relationships with significant positive slopes (12) resulting was not significantly different from the number of those with significant negative slopes (5) according to the chi-square test ($\chi^2$ = 2.882, P = 0.089), showing independence of the $W_r$ values calculated by means of the EmP $W_s$ equation developed by the size of the fish.

Applying the EmpQ method, the value of the slope was not significantly different from zero for both terms of the equation ($P_{quadratic}$ = 0.508, $P_{linear}$ = 0.763), indicating that the EmP $W_s$ equation was not influenced by fish length (Figure 4).





**Figure 3.** Relationship between variance/mean for $\log_{10}$ of weight (W) and total length at 10-mm intervals for the determination of the minimum total length of *Squalius pursakensis*. The dotted line indicates the value of 0.01.

## 4. Discussion

In this study, a standard weight equation was developed for *S. pursakensis*, an endemic species of the Sakarya and Porsuk drainages in Northwest Anatolia (Turkey). Moreover, a new maximum total length (445 mm) for this species was found from Seydi Suyu in Sakarya Basin. Therefore, it can be argued that the representativeness of the dataset used in the present study has contributed to strengthen the validity of the findings. Maximum total length was previously reported as 392 mm from Sarıyar Reservoir by Ekmekçi (1996). Although it is well known that individuals in populations exposed to high levels of fishery pressure reach relatively smaller lengths (Emiroğlu et al., 2012), this is not the case for *S. pursakensis*, as there is no high fishery pressure on the species other than the insignificant effect of local anglers in its distribution range. The finding of large specimens of a species could be incidental or attributed to inadequate current knowledge on the species. However, given that congeneric species of *S. pursakensis* may reach 600 mm SL (*S. cephalus*) (Kottelat and Freyhof, 2007), it can grow above the maximum length found in the present study.

Relative weight is currently widely used to perform condition analysis of many species, but its applicability is often limited by the lack of species-specific standard weight equations. With regard to Turkey, the only $W_s$ equations available are those for European perch *Perca fluviatilis* (Giannetto et al., 2012a) and Aegean chub *Squalius fellowesii* (Giannetto et al., 2012c).

Previous studies on *S. pursakensis*, which was cited mainly as *Leuciscus cephalus*, focused on some basic reproduction and growth features, including condition indices (Erk'akan, 1981; Ekmekçi, 1996; Emiroğlu et al., 2001; Bostancı and Polat, 2009; Innal, 2010). Interestingly, most of the previous studies were conducted in artificial water bodies (reservoirs and ponds) (e.g., Ekmekçi, 1996; Bostancı and Polat, 2009; Yiğit et al., 2008; Innal, 2010) with the exception of 2 studies, which were from the Sakarya (Erk'akan, 1981) and Porsuk (Emiroğlu et al., 2001) rivers. Yiğit et al. (2008) analyzed different condition indices with the aim of finding the best-fitting condition factor for *S. pursakensis* and found that the one based on the height of fish along with length and weight proposed by Jones et al. (1999) was the best-fitting condition factor. However, this factor is not useful for broad comparisons due to the lack

**Table.** Number of populations and individuals for each length class of 10 mm in the development dataset of *Squalius pursakensis*. The length classes in bold are those removed from the dataset because they had fewer than 3 populations.

| TL (mm) | Populations | Individuals | TL (mm) | Populations | Individuals |
|---|---|---|---|---|---|
| 80 | 31 | 62 | 270 | 15 | 18 |
| 90 | 35 | 70 | 280 | 13 | 19 |
| 100 | 42 | 105 | 290 | 7 | 10 |
| 110 | 52 | 153 | 300 | 6 | 9 |
| 120 | 67 | 203 | 310 | 7 | 10 |
| 130 | 63 | 209 | 320 | 3 | 5 |
| 140 | 70 | 257 | 330 | 3 | 5 |
| 150 | 59 | 189 | 340 | 3 | 4 |
| 160 | 62 | 184 | **350** | **1** | **1** |
| 170 | 54 | 133 | **360** | - | - |
| 180 | 49 | 121 | **370** | **3** | **3** |
| 190 | 41 | 93 | **380** | **2** | **3** |
| 200 | 40 | 71 | **390** | - | - |
| 210 | 39 | 60 | **400** | - | - |
| 220 | 38 | 68 | **410** | - | - |
| 230 | 26 | 41 | **420** | - | - |
| 240 | 20 | 26 | **430** | - | - |
| 250 | 24 | 27 | **440** | - | - |
| 260 | 21 | 24 | **450** | **1** | **1** |





**Figure 4.** Plots showing the distribution of the residuals (a) and the results of the application of the empirical quartiles (EmpQ) method (b) used to investigate potential length bias in the standard mass ($W_s$) equation for *Squalius pursakensis*. Residuals = standardized residuals of the regression; fitted values = values obtained by the model fit; standardized 75th percentile mean W = standardized 75th percentile mean weights calculated by $W_s$ equation.

of available data (i.e. height is usually not measured) and its uncommon use (i.e. this condition factor should be site-specific). Indeed, this was the case for Yiğit et al. (2008), as they could use only the Fulton condition factor (*K*) for comparisons with other studies, though it had the lowest coefficient factor among all analyzed condition indices. In this context, relative weight is the best alternative to condition factors commonly used in fishery biology, because it provides a single species-specific equation for the species, which allows reliable comparisons among different locations and specimens of different lengths.

Further research is encouraged to extend the use of this methodology to other species, with particular attention to those that are endemic. Turkey is an important area for fish biodiversity, being characterized by a high number of endemic species (78 species). Notably, almost half of the endemic species in Turkey are classified as 'Critically Endangered' and 32% as 'Endangered' (Fricke et al., 2007). These species often have a very small area of distribution (like *S. pursakensis*), are rarely studied, and currently are threatened by the presence of an increasing number of nonnative species introduced by stocking practices that have become very common in Turkish waters to increase fish production and sport fishing (Innal and Erk'akan, 2006; Aydın et al., 2011; Önsoy et al., 2011; Tarkan et al., 2012). For these reasons, all management tools that can assist in conserving the populations of these endemic species, such as *S. pursakensis*, would be advantageous in assessing the population-level responses to ecosystem disturbance.

The relative weight, together with other population metrics (e.g., age and growth), could allow researchers to increase basic knowledge of the population ecology of *S. pursakensis*, which will be useful in improving its management and conservation status (Murphy et al., 1991; Blackwell et al., 2000). Low values of relative weight will permit, by comparison of data of different periods, detection of any decline in the condition of specimens, probably due to environmental alterations or biological disturbance (e.g., presence of nonnative species) (Giannetto et al., 2012b). The results of previous studies reported clear habitat-based variations in age and growth features for other chub species, such as Aegean chub (Balık et al., 2004; Dirican and Barlas, 2007). These variations are more prominent between populations inhabiting lentic and lotic areas (Şaşı and Balık, 2003; Koç et al., 2007). Because of the positive correlation between fish growth and environmental quality (Bister et al., 2000), relative weight could be an easy and powerful tool for identifying ecological changes, such as the incidence of phenomena of intraspecies competition (Johnson, 1992), impact of nonnative species Giannetto et al., 2012b), or environmental disturbances (Gabelhouse, 1991; Hubert et al., 1994; Liao et al., 1995). Considering that *S. pursakensis* is commonly found in reservoirs, which are considered as unfavorable and variable environments for fish species (Tarkan, 2007; Emiroğlu et al., 2012), monitoring ecological changes via simple and effective tools such as relative weight is useful.

The EmP $W_s$ equation developed for *S. pursakensis* is not biased by fish length according to all methods used for the validation (Willis, EmpQ, and analysis of the residuals of the equation). The results highlighted the reliability of the EmP method in developing standard weight equations, confirming those obtained by previous studies (Rennie and Verdon, 2008; Ogle and Winfield, 2009; Giannetto et al., 2011, 2012a, 2012c; Lorenzoni et al., 2012). On the basis of these results, the use of the EmP equation to determine $W_r$ for *S. pursakensis* throughout its area of distribution is suggested.

**Acknowledgments**

This study was jointly funded by the Anadolu University Research Fund (Project No. 1101F011) and the Osmangazi University Research Fund (Project No. 2012/19A113).

The statistical analyses of data of this research were conducted within the 2216 "Research Fellowship Programme for International Researchers" supported by the Scientific and Technological Research Council of Turkey (TÜBİTAK).

## References

Aydın H, Gaygusuz Ö, Tarkan AS, Top N, Emiroğlu Ö, Gürsoy Gaygusuz Ç (2011). Invasion of freshwater bodies in the Marmara region (northwestern Turkey) by nonnative gibel carp *Carassius gibelio* (Bloch, 1782). Turk J Zool 35: 829–836.

Anderson RO, Neumann RM (1996). Length, weight, and associated structural indices. In: Murphy BR, Willis DR, editors. Fisheries Techniques. 2nd ed. Bethesda, MD, USA: American Fisheries Society, pp. 447–481.

Balık S, Sarı HM, Ustaoğlu MR, İlhan A (2004). Age and growth characteristics of chub (*Leuciscus cephalus* L., 1758) population in Işıklı Lake, Çivril, Denizli, Turkey. E.U. Journal of Fisheries & Aquatic Sciences 21: 257–262 (article in Turkish with English abstract).

Begg GA, Friedland KD, Pearce JB (1999). Stock identification and its role in stock assessment and fisheries management: an overview. Fish Res 43: 1–8.

Bister TJ, Willis DW, Brown ML, Jordan SM, Neumann RM, Quist MC, Guy CS (2000). Proposed standard weight ($W_s$) equations and standard length categories for 18 warmwater nongame and riverine fish species. North Am J Fish Manage 20: 570–574.

Blackwell BG, Brown ML, Willis DW (2000). Relative weight ($W_r$) status and current use in fisheries assessment and management. Rev Fish Sci 8: 1–44.

Bostancı D, Polat N (2009). Age determination and some population characteristics of chub (*Squalius cephalus* L., 1758) in the Çamlıdere Dam Lake (Ankara, Turkey). Turkish Journal of Science & Technology 4: 25–30.

Dirican S, Barlas M (2007). A study on the condition factors of *Leuciscus cephalus* (L., 1758) and *Barbus capito pectoralis* Heckel, 1843 populations living in Dipsiz-Çine (Muğla-Aydın) Stream. Afyon Kocatepe University Journal of Science 7: 105–114 (article in Turkish with English abstract).

Durand JD, Ünlü E, Doadrio I, Pipoyan S, Templeton AR (2000). Origin, radiation, dispersion and allopatric hybridization in the chub *Leuciscus cephalus*. Proc Biol Sci 267: 1687–1697.

Ekmekçi FG (1996). Sarıyar Baraj Gölü'nde yaşayan tatlısu kefalinin (*Leuciscus cephalus* Linnaeus, 1758) büyüme ve üreme özellikleri. Turk J Zool 20: 95–106 (article in Turkish).

Emiroğlu Ö, Tarkan AS, Top N, Başkurt S, Sülün Ş (2012). Growth and life history traits of a highly exploited population of non-native gibel carp, *Carassius gibelio* from a large eutrophic lake (Lake Uluabat, NW Turkey): is reproduction the key factor for establishment success? Turk J Fish Aquat Sci 12: 925–936.

Emiroğlu Ö, Yetim M, Kuyumcu N (2001). Porsuk Çayı havzasında yaşayan tatlı su Kefali'nin (*Leuciscus cephalus* L.) büyüme performansının araştırılması. In: XI Ulusal Su Ürünleri Sempozyumu, September 2001. Hatay, Turkey: Özet Kitabı Cilt, pp. 11–25 (symposium proceeding in Turkish).

Fechhelm RG, Griffiths WB, Wilson WJ, Gallaway BJ, Bryan JD (1995). Intra- and interseasonal changes in the relative condition and proximate body composition of broad whitefish from the Prudhoe Bay Region of Alaska. Trans Am Fish Soc 124: 508–519.

Fricke R, Bilecenoğlu M, Sarı HM (2007). Annotated checklist of fish and lamprey species (Gnathostomata and Petromyzontomorphi) of Turkey, including a Red List of threatened and declining species. Stuttg Beitr Natkd A Biol 706: 1–169.

Froese R (2006). Cube law, condition factor and weight-length relationships: history, meta-analysis and recommendations. J Appl Ichthyol 22: 241–253.

Fulton TW (1911). The Sovereignty of the Sea: An Historical Account of the Claims of England to the Dominion of the British Seas and of the Evolution of the Territorial Waters. Edinburgh, UK: William Blackwood.

Gabelhouse DW (1991). Seasonal changes in body condition of white crappies and relations to length and growth in Melvem Reservoir, Kansas. North Am J Fish Manage 11: 50–56.

Gerow KG (2010). Biases with the regression line percentile method and the fallacy of a single standard weight. North Am J Fish Manage 30: 679–690.

Gerow KG, Anderson-Sprecher R, Hubert WA (2005). A new method to compute standard-weight equations that reduce length-related bias. North Am J Fish Manage 25: 1288–1300.

Gerow KG, Hubert WA, Anderson-Sprecher R (2004). An alternative approach to detection of length-related biases in standard-weight equations. North Am J Fish Manage 24: 903–910.

Giannetto D, Carosi A, Franchi E, La Porta G, Lorenzoni M (2012a). Proposed standard weight ($W_s$) equation for European perch *Perca fluviatilis* Linnaeus, 1758. J Appl Ichthyol 28: 34–39.

Giannetto D, Carosi A, Franchi E, Pedicillo G, Pompei L, Lorenzoni M (2012b). Assessing the impact of non-native freshwater fishes on native species using relative weight. Knowl Manag Aquat Ec 404: 03.

Giannetto D, La Porta G, Maio G, Pizzul E, Turin P, Lorenzoni M (2011). Proposed standard mass equations for European chub *Leuciscus cephalus* in Italy. J Fish Bio 78: 1890–1899.

Giannetto D, Pompei L, Lorenzoni M, Tarkan AS (2012c). Empirical standard weight equation for Aegean Chub *Squalius fellowesii*, an endemic freshwater fish species of Western Anatolia. North Am J Fish Manage 32: 1102–1107.

Hubert WA, Gipson RD, Whaley RA (1994). Interpreting relative weight values of lake trout stocks. North Am J Fish Manage 14: 212–215.

Innal D (2010). Population structures and some growth properties of three cyprinid species [*Squalius cephalus* (Linnaeus, 1758); *Tinca tinca* (Linnaeus, 1758) and *Alburnus escherichii* Steindachner, 1897] living in Camkoru Pond (Ankara-Turkey). Kafkas Univ Vet Fak Derg 16: 297–304.

Innal D, Erk'akan F (2006). Effect of exotic and translocated fish species in the inland waters of Turkey. Rev Fish Biol Fisher 16: 39–50.

Johnson SL, Rahel FJ, Hubert WA (1992). Factors influencing the size structure of brook trout populations in beaver ponds in Wyoming. North Am J Fish Manage 12: 118–124.

Jones RE, Petrell RJ, Pauly D (1999). Using modified length-weight relationships to assess the condition of fish. Aquacul Eng 20: 261–276.

Koç HT, Erdoğan Z, Tinkci M, Treer T (2007). Age, growth and reproductive characteristics of chub, (*Leuciscus cephalus* L., 1758) in the Ikizcetepeler Dam lake (Balikesir), Turkey. J Appl Ichthyol 23:19–24.

Kottelat M, Freyhof J (2007). Handbook of European Freshwater Fishes. Cornol, Switzerland: Publications Kottelat.

Le Cren ED (1951). The length-weight relationship and seasonal cycle in gonad weight and condition in the perch (*Perca fluviatilis*). J Anim Ecol 20: 201–219.

Liao H, Pierce CL, Wahl DH, Rasmussen JB, Leggett WC (1995). Relative weight ($W_r$) as a field assessment tool: relationships with growth, prey biomass, and environmental conditions. Trans Am Fish Soc 124: 387–400.

Lorenzoni M, Giannetto D, Maio G, Pizzul E, Pompei L, Turin P, Vincenzi S, Crivelli A (2012). Empirical standard mass equation for *Salmo marmoratus*. J Fish Bio 81: 2086–2091.

Murphy BR, Brown ML, Springer TA (1990). Evaluation of the relative weight ($W_r$) index with new applications to walleye. North Am J Fish Manage 10: 85–97.

Murphy BR, Willis DW, Springer TA (1991). The relative weight index in fisheries management: status and needs. Fisheries 16: 30–38.

Ogle DH, Winfield IJ (2009). Ruffe length-weight relationships with a proposed standard weight equation. North Am J Fish Manage 29: 850–858.

Önsoy B, Filiz H, Tarkan AS, Bilge G, Tarkan AN (2011). Occurrence of non-native fishes in a small man-made lake (Lake Ula, Muğla): past, present, future perspectives. Turk J Fish Aquat Sci 11: 209–215.

Özulug M, Freyhof J (2011). Revision of the genus *Squalius* in Western and Central Anatolia, with description of four new species (Teleostei: Cyprinidae). Ichthyol Explor Fres 22: 167.

Rennie MD, Verdon R (2008). Development and evaluation of condition indices for the lake whitefish. North Am J Fish Manage 28: 1270–1293.

Rypel A, Richter TJ (2008). Empirical percentile standard weight equation for the blacktail redhorse. North Am J Fish Manage 28: 1843–1846.

Şaşı H, Balık S (2003). Age, growth and sex ratio of chub (*Leuciscus cephalus* L., 1758) in Topçam Dam Lake (Aydın). E.U. Journal of Fisheries & Aquatic Sciences 20: 503–515.

Tarkan AN, Gaygusuz Ö, Tarkan AS, Gürsoy Ç, Acıpınar H (2007). Interannual variability of fecundity and egg size of an invasive cyprinid, *Carassius gibelio*: effects of density-dependent and density-independent factors. J Freshwater Ecol 22: 11–17.

Tarkan AS, Copp GH, Top N, Özdemir N, Önsoy B, Bilge G, Filiz H, Yapıcı S, Ekmekçi G, Kırankaya Ş et al. (2012). Are introduced gibel carp *Carassius gibelio* in Turkey more invasive in artificial than in natural waters? Fish Manage Ecol 19: 178–187.

Wege GJ, Anderson RO (1978). Relative weight ($W_r$): a new index of condition for largemouth bass. In: Novinger GD, Dillard JG, editors. New Approaches to the Management of Small Impoundments. Bethesda, MD, USA: Special Publications, American Fisheries Society, pp. 79–91.

Willis DW, Guy CS, Murphy BR (1991). Development and evaluation of a standard weight ($W_s$) equation for yellow perch. North Am J Fish Manage 11: 374–380.

Yiğit S, Ergönül MB, Altındağ A (2008). The growth features of chub *Squalius cephalus* and comparison of five different condition indices. Cybium 32: 317–319.